


Claims

1. A compound having the formula



5

a pharmaceutically acceptable acid addition salt thereof, or a stereochemically isomeric form thereof, wherein

Alk is C₁₋₄alkanediyl;

R¹ is hydrogen, C₁₋₄alkyl or halo;

10

R² is C₁₋₄alkyl;

R³ is hydroxy or R⁴-C(=O)O-; and

R⁴ is C₁₋₁₉alkyl.

15

2. A compound according to claim 1 wherein R³ is substituted on the 9 position of the 6,7,8,9-tetrahydro-2-C₁₋₄alkyl-4H-pyrido[1,2-a]pyrimidin-4-one moiety.

3. A compound according to claim 2 wherein Alk is ethanediyl, R¹ is halo and R² is methyl.

20

4. A compound according to claim 3 wherein R¹ is 6-fluoro.


5. A compound according to any of claims 1 to 4 wherein R⁴ is heptyl, nonyl, undecyl or tridecyl.

25

6. A compound according to claim 1 wherein the compound is selected from 3-[2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)-1-piperidinyl]ethyl]-6,7,8,9-tetrahydro-9-hydroxy-2-methyl-4H-pyrido[1,2-a]pyrimidin-4-one, a pharmaceutically acceptable acid addition salt thereof, or an enantiomeric form thereof.

30

7. An antipsychotic composition comprising an inert carrier and as active ingredient an antipsychotic effective amount of a compound having the formula



a pharmaceutically acceptable acid addition salt thereof, or a stereochemically isomeric form thereof, wherein

- 5 Alk is C₁₋₄alkanediyl;
 R¹ is hydrogen, C₁₋₄alkyl or halo;
 R² is C₁₋₄alkyl;
 R³ is hydroxy or R⁴-C(=O)O-; and
 R⁴ is C₁₋₁₉alkyl.

10

8. A composition according to claim 7 wherein R³ is substituted on the 9 position of the 6,7,8,9-tetrahydro-2-C₁₋₄alkyl-4H-pyrido[1,2-a]pyrimidin-4-one moiety.

15

9. A composition according to claim 8 wherein Alk is ethanediyl, R¹ is halo and R² is methyl.

10. A composition according to claim 9 wherein R¹ is 6-fluoro.

20


11. A composition according to any of claims 7 to 10 wherein R⁴ is heptyl, nonyl, undecyl or tridecyl.

25

12. A composition according to claim 7 wherein the compound is selected from 3-[2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)-1-piperidinyl]ethyl]-6,7,8,9-tetrahydro-9-hydroxy-2-methyl-4H-pyrido[1,2-a]pyrimidin-4-one, a pharmaceutically acceptable acid addition salt thereof, or an enantiomeric form thereof.

30

13. A method of treating warm-blooded animals suffering from psychotic diseases, which method comprises the administration to said warm-blooded animals of an antipsychotic effective amount of a chemical compound having the formula



a pharmaceutically acceptable acid addition salt thereof, or a stereochemically isomeric form thereof, wherein

- 5 Alk is C₁-4alkanediyl;
 R¹ is hydrogen, C₁-4alkyl or halo;
 R² is C₁-4alkyl;
 R³ is hydroxy or R⁴-C(=O)O-; and
 R⁴ is C₁-19alkyl.

10

14. A method according to claim 13 wherein R³ is substituted on the 9 position of the 6,7,8,9-tetrahydro-2-C₁-4alkyl-4H-pyrido[1,2-a]pyrimidin-4-one moiety.

15

15. A method according to claim 14 wherein Alk is ethanediyl, R¹ is halo and R² is methyl.

16. A method according to claim 15 wherein R¹ is 6-fluoro.


20

17. A method according to any of claims 13 to 16 wherein R⁴ is heptyl, nonyl, undecyl or tridecyl.

25

18. A method according to claim 13 wherein the compound is selected from 3-[2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)-1-piperidinyl]ethyl]-6,7,8,9-tetrahydro-9-hydroxy-2-methyl-4H-pyrido[1,2-a]pyrimidin-4-one, a pharmaceutically acceptable acid addition salt thereof, or an enantiomeric form thereof.


19. A process for preparing a compound having the formula



a pharmaceutically acceptable acid addition salt thereof, or a stereochemically isomeric form thereof, wherein


- 5 Alk is C₁-4alkanediyl;
 R¹ is hydrogen, C₁-4alkyl or halo;
 R² is C₁-4alkyl;
 R³ is hydroxy or R⁴-C(=O)O-; and
 R⁴ is C₁-19alkyl; characterized by

- 10 a) N-alkylating a 3-piperidinyl-1,2-benzisoxazole of formula



wherein R¹ is as defined under formula (I) with an alkylating reagent of formula


15



wherein R², R³ and Alk are as defined under formula (I) and W represents a leaving group in a reaction-inert solvent at an elevated temperature;

20


- b) cyclizing an oxime of formula



wherein R^1 , R^2 , R^3 and Alk are as defined under formula (I) and Y is a reactive leaving group in a reaction-inert solvent at an elevated temperature and in the presence of a base;


5

c) cyclizing an activated oxime derivative




10 wherein R^1 , R^2 , R^3 and Alk are as defined under formula (I) and L is an acid residue in a reaction-inert solvent at an elevated temperature and in the presence of a base;

d) O-acylating a compound of formula



15


wherein R^1 , R^2 and Alk are as defined under formula (I) with a carboxylic acid of formula




20

or a reactive functional derivative thereof, wherein R^4 is as defined under formula (I) in a reaction-inert solvent;

e) cyclizing an amidine of formula




5 wherein R³ is as defined under formula (I) with a β-dicarbonyl intermediate of formula




10 wherein R¹, R² and Alk are as defined under formula (I) and R⁵ is a leaving group in a reaction-inert solvent at an elevated temperature;

f) cyclizing a reagent of formula




15 wherein R³ is as defined under formula (I) and R⁵ is a leaving group with an enamine of formula



20 wherein R¹, R² and Alk are as defined under formula (I) and R⁵ is a leaving group, in a reaction-inert solvent at an elevated temperature; or

25 g) preparing the enantiomeric forms of the compounds of formula



- wherein R^1 , R^2 and Alk are as defined under formula (I), by converting the racemic mixtures of the compounds of formula (I-a) with a resolving agent to a mixture of
- 5 diastereomeric salts or compounds, physically separating said mixture of diastereomeric salts or compounds and converting said separated diastereomeric salts or compounds into the corresponding enantiomeric forms of the compounds of formula (I-a);
- and if desired, converting the compounds of formula (I) into a therapeutically active non-
- 10 toxic acid addition salt form by treatment with an acid; or conversely, converting the acid salt into the free base with alkali;

Inv.
B2